

A SUB-DENSITY THEOREM OF STURM-LIOUVILLE EIGENVALUE PROBLEM WITH FINITELY MANY SINGULARITIES

LUNG-HUI CHEN

National Chung Cheng University, Min-Hsiung, Taiwan
E-mail: mr.lunghuichen@gmail.com

Abstract. We study the distribution of the Sturm-Liouville eigenvalues of a potential with finitely many singularities. There is an asymptotically periodical structure on this class of eigenvalues as described by the entire function theory. We describe the singularities of its potential function explicitly in its eigenvalue asymptotics.

MSC2010 numbers: 34B24, 35P25, 35R30.

Keywords: Sturm-Liouville problem; singular eigenvalue problem; Wilder's theorem.

1. INTRODUCTION AND MAIN RESULT

In this short note, we study the eigenvalue distribution for the following differential equation.

$$(1.1) \quad \begin{cases} -y''(x) + p(x)y(x) = \omega^2 y(x), & 0 < x < \pi; \\ y(0; \omega) = 0, \quad y'(0; \omega) = 1; \\ y(\pi; \omega) = 0, \end{cases}$$

where

$$(1.2) \quad p(x) = \sum_{m=0}^M p_m(x) + r_M(x)$$

with $r_M(x) \in C^M[0, \pi]$; Most importantly,

$$(1.3) \quad p_m(x) = \sum_k c_{m,k} l_{[x_{m,k}, \infty)}(x) (x - x_{m,k})^m / m!, \quad m \geq 1;$$

$$(1.4) \quad p_0(x) = \sum_k c_{0,k} l_{[x_{0,k}, \infty)}(x),$$

where $\{x_{m,k}\}_{m,k} \in (0, \frac{\pi}{2})$ and $\{c_{m,k}\}_{m,k} \in \mathbb{R}$. We are dealing with a piecewise $C^M[0, \pi]$ potential function $p(x)$. For each m , $x_{m,k}$ are distinct and $p(x)$ has a jump at m -th derivative at $x_{m,k}$. We assume nontrivially the $\{x_{m,k}\}_{m,k} \in (0, \frac{\pi}{2})$ has J elements and are all distinct. If there are two singular points symmetrically to the middle point located in $(0, \pi)$, then our method doesn't apply in this case.

It is asked by Carlson, Threadgill and Shubin [1]: How are the singularities of p manifested in the distribution of eigenvalues? Being considered as a function of ω , $y(\pi; \omega)$ is an entire function of ω . Moreover, the zeros of $y(\pi; \omega)$ are the Dirichlet eigenvalues of the system (1.1). To study the asymptotics of Dirichlet eigenvalues, we examine the zeros of entire function $y(\pi; \omega)$. We try to answer the question from the point of view of complex analysis in this particular setting. In [1], a distribution of the eigenvalues with coefficients in terms of spectral invariants is described in [1, Theorem 4.4] applying the Newton's method. In this paper, we try to characterize the distribution of the eigenvalues explicitly in terms of the singularities themselves and find the composites of the Dirichlet eigenvalues. Can one really hear the singularities of the potential p ? The next statement is the main result of this paper.

Theorem 1.1. *Let $\{\omega_j\}_{j=1}^J$ be the rearrangement of the singular points $\{x_{m,k}\}_{m,k}$ such that $0 =: \omega_0 < \omega_1 < \omega_2 < \dots < \omega_J < \omega_{J+1} := \frac{\pi}{2}$. There exist exactly $2J + 2$ subsequences of the zeros of $y(\pi; \omega)$, denoted as $\{z_{n_l}\}$, where $l = 1, 2, \dots, 2J + 2$, such that*

$$(1.5) \quad z_{n_l} \sim \frac{n_l \pi}{\omega_l - \omega_{l-1}} + O(1), \text{ as } n_l \rightarrow \pm\infty \text{ in } \mathbb{Z};$$

$$(1.6) \quad \bigcup_{l=1}^{2J+2} \{z_{n_l}\} = \{z_n\},$$

in which $\{z_n\}$ are the zeros of $y(\pi; \omega)$.

In particular, we recover the point set $\{\omega_j\}_{j=1}^J$ from the subsequences of Dirichlet eigenvalues corresponding to each of these points. We may refine the asymptotics (1.1) to next order by the method in [9, p. 37]:

$$(1.7) \quad z_{n_l} \sim \frac{n_l \pi}{\omega_l - \omega_{l-1}} + O\left(\frac{1}{n_l}\right), \text{ as } n_l \rightarrow \pm\infty \text{ in } \mathbb{Z}.$$

This is the only eigenvalue asymptotics containing the information on the position of the singularities of a given potential function known to the author. We may compare the result in [6, 7, 9]. However, in [6], they considered a much general class of potential functions. One may sum up all of the subsequences to obtain the classic eigenvalue density as in [7, 9].

We start with the asymptotic expansion of the solution of (1.1) which we refer to [1, 2]. The following asymptotics holds:

$$(1.8) \quad y(\pi; \omega) = \frac{\sin\{\omega\pi\}}{\omega} - \frac{1}{2} \frac{\cos\{\omega\pi\}}{\omega^2} \int_0^\pi p(t) dt + \\ + 2 \sum_{m=1}^{M-1} (-1)^m [2\omega]^{-2m} \cos\{\omega\pi\} P_{m-1}(\pi) + 2 \sum_{m=1}^{M-1} (-1)^{m+1} [2\omega]^{-2m-1} \sin\{\omega\pi\} Q_{m-1}(\pi)$$

$$\begin{aligned}
& +2 \sum_{m=0}^{\frac{M-1}{2}-1} (-1)^m [2\omega]^{-2m-3} \times \left\{ \sum_{x_{2m,k} < \pi} c_{2m,k} \sin\{\omega[x - 2x_{2m,k}]\} \right\} \\
& +2 \sum_{m=0}^{\frac{M-1}{2}-1} (-1)^{m+1} [2\omega]^{-2m-4} \times \left\{ \sum_{x_{2m+1,k} < \pi} c_{2m+1,k} \cos\{\omega[x - 2x_{2m+1,k}]\} \right\} + O\left(\frac{\exp \omega \pi}{\omega^M}\right),
\end{aligned}$$

if $\omega \in \mathbb{C}$ and

$$\begin{aligned}
P_0(x) &:= \int_0^x p(t_1) dt_1; \quad P_m(x) := q_{2m-1}(p; x) - q_{2m-1}(p; 0), \quad m \in \mathbb{N}; \\
Q_m(x) &:= q_{2m}(p; x) + q_{2m}(p; 0), \quad m \in \mathbb{N}_0; \\
q_m(p; x) &:= \left[\sum_{k=m}^M p_k(x) + r_M(x) \right]^{(m)}, \quad r_M(x) \in C^M[0, \pi].
\end{aligned}$$

This is essentially the (3.e) in [1, p. 84]. However, we deal with $\omega \in \mathbb{C}$ in this paper. The only difference is in the big O -term in the end of (1.8). We refer the proof to [1, p. 84], and also [9], which comes from the repeated integration by parts. We will apply the Wilder's theorem to (1.8) which is a sum of asymptotically hyperbolic series to obtain the asymptotics of the Dirichlet eigenvalues.

2. THE WILDER'S THEOREM

There is an asymptotic periodic structure [4, 5, 8] within the zero set of the asymptotically hyperbolic sum, say, the asymptotic expansion (1.8). We refer to [5, 8] for a comprehensive study on the zero distribution theory of this kind. To be more convincing, we start with the following theorem. One can bypass this part if familiar with the entire function theory. The indexing in this section is independent of the others.

Theorem 2.1 (Dickson [4]). *Let*

$$(2.1) \quad R(\alpha, s, h) := \{z = x + iy \in \mathbb{C} \mid |x| \leq h, y \in [\alpha, \alpha + s]\};$$

$$(2.2) \quad N_g(R(\alpha, s, h)) := \{ \text{the number of zeros of } g(z) \text{ in } R(\alpha, s, h) \},$$

in which

$$g(z) = \sum_{j=1}^n A_j e^{\omega_j z},$$

where $z = x + iy$, $A_j \neq 0$, $\omega_1 < \omega_2 < \dots < \omega_n$. Then, there exists $K > 0$ such that

- (1) each zero of g is in $|x| < K$;
- (2) for each pair of reals (α, s) with $s > 0$,

$$(2.3) \quad |N_g(R(\alpha, s, K)) - s(\omega_n - \omega_1)/(2\pi)| \leq n - 1.$$

Let us acquire a more sophisticated theorem of this type. Let

$$(2.4) \quad f(z) = \sum_{j=1}^n A_j z^{m_j} [1 + \epsilon(z)] e^{\omega_j z},$$

where $n > 1$ and A_j and ω_j are complex numbers such that $A_j \neq 0$ and the ω_j are distinct; the m_j are non-negative integers; the functions ϵ are analytic for $|z| \geq r_0 \geq 0$ with $\lim_{z \rightarrow \infty} \epsilon(z) = 0$. When we are talking about the zeros of $f(z)$, we are referring to its zeros outside certain open ball around the origin.

We set up the following quantities to the $f(z)$ in (2.4): Let Q be the broken line given by the $\bar{\omega}_j$ given in (2.4) with $\bar{\omega}_1, \dots, \bar{\omega}_\sigma$ as its vertices. The indices are labeled counterclockwise. Let L_k be the line segment $[\bar{\omega}_k, \bar{\omega}_{k+1}]$ and $\phi_k := \arg\{\bar{\omega}_k - \bar{\omega}_{k+1}\}$ in $[-\frac{\pi}{2}, \frac{3\pi}{2}]$. Let

$$(2.5) \quad e_k = e^{i\phi_k}.$$

Certain $\bar{\omega}_p$ on L_k are assigned doubly indexed subscripts as follows: Let the convex hull of $\bar{\omega}_k, \bar{\omega}_{k+1}$ and $\tau_p = \bar{\omega}_p + im_p e_k$ in which $\bar{\omega}_p$ on L_k ; assign subscripts $j = 1, \dots, \sigma_k$ to ω_{kj} so that $\omega_{k1} = \omega_k, \omega_{k\sigma_k} = \omega_{k+1}$ and τ_{kj} are vertices of this convex hull and preceding in a counterclockwise direction from $\bar{\omega}_k + im_k e_k$ to $\bar{\omega}_{k+1} + im_{k+1} e_k$. For $j = 1, \dots, \sigma_k - 1$,

$$(2.6) \quad L_{kj} := [\tau_{kj}, \tau_{kj+1}]; \quad \mu_{kj} := \frac{m_{kj} - m_{kj+1}}{(\omega_{kj} - \omega_{kj+1})e_k}$$

which is real; n_{kj} is the number of τ_p on L_{kj} . In particular, if L_{kj} in an interval with exactly two end points, then we have $n_{kj} = 2$.

Moreover, for $j = 1, \dots, \sigma_k - 1$ and $h > 0$, we define

$$(2.7) \quad V_{kj}(h) := \{z | \Im(z/e_k) \geq 0, |\Re(z/e_k) + \mu_{kj} \log |z|| \leq h\}.$$

$T_k(\theta)$ is defined to be a closed sector with vertex at zero of opening 2θ about the outward normal to L_k through the origin. For the same k and j and each triple of reals (α, s, h) , $s > 0$ and $h > 0$, the set

$$(2.8) \quad R_{kj}(\alpha, s, h) := \{z | \Im(z/e_k) + \mu_{kj} \arg z \in [\alpha, \alpha + s], |\Re(z/e_k) + \mu_{kj} \log |z|| \leq h\},$$

where $\arg z \in (\phi_k, \phi_k + \pi)$ and $R_{kj}(\alpha, s, h)$ is in $V_{kj}(h) \cap U'_k(\theta)$. They are asymptotically logarithmic tubular neighborhoods. We refer to [4] for a comprehensive study. Now we state the following theorem.

Theorem 2.2 (Dickson [4]). *Let $f(z)$ be given as in (2.4). Then, there exists $h > 0$ such that*

- (1) *all but a finite number of zeros of f of modulus greater than r_0 are in*
 $\bigcup_{k,j} V_{kj};$

(2) for each pair of positive reals ϵ and s_0 , there exists an $\alpha_0 = \alpha_0(\epsilon, s_0)$ such that whenever $\alpha \geq \alpha_0$ and $s \geq s_0$,

$$(2.9) \quad |N_f(R_{kj}(\alpha, s, h)) - s|\omega_{kj+1} - \omega_{kj}|/(2\pi)| < n_{kj} - 1 + \epsilon.$$

This is exactly stated as in [4]. The proof is in [5, Theorem 2, p.21]. We refer to [3] for another application of this theorem.

3. PROOF OF THEOREM 1.1

To apply Theorem 2.2 to the hyperbolic sum (1.8), we rewrite (1.8): It is well-known [9] that there is a C_δ depending on the distance to the zeros of $\sin \omega\pi$ such that

$$(3.1) \quad \exp |\Im \omega \pi| < C_\delta \sin \{\omega \pi\}.$$

Hence, (1.8) becomes

$$\begin{aligned} y(\pi; \omega) &= \frac{\sin \{\omega \pi\}}{\omega} [1 + O(\frac{C_\delta}{\omega^{M-1}})] - \frac{1}{2} \frac{\cos \{\omega \pi\}}{\omega^2} \int_0^\pi p(t) dt \\ &+ 2 \sum_{m=1}^{\frac{M-1}{2}} (-1)^m [2\omega]^{-2m} \cos \{\omega \pi\} P_{m-1}(\pi) + 2 \sum_{m=1}^{\frac{M-1}{2}} (-1)^{m+1} [2\omega]^{-2m-1} \sin \{\omega \pi\} Q_{m-1}(\pi) \\ &+ 2 \sum_{m=0}^{\frac{M-1}{2}-1} (-1)^m [2\omega]^{-2m-3} \times \left\{ \sum_{x_{2m,k} < \pi} c_{2m,k} \sin \{\omega[x - 2x_{2m,k}]\} \right\} \\ &+ 2 \sum_{m=0}^{\frac{M-1}{2}-1} (-1)^{m+1} [2\omega]^{-2m-4} \times \left\{ \sum_{x_{2m+1,k} < \pi} c_{2m+1,k} \cos \{\omega[x - 2x_{2m+1,k}]\} \right\}, \end{aligned}$$

$\omega \notin \mathbf{Z}$. Now we rearrange according to their exponential powers by the theorem assumption to the following form:

$$\begin{aligned} y(\pi; \omega) &= \left\{ \frac{-1}{2i\omega} [1 + O(\frac{C_\delta}{\omega^{M-1}})] + \sum_{m=1}^{\frac{M-1}{2}} (-1)^m [2\omega]^{-2m} P_{m-1}(\pi) \right. \\ &+ i \sum_{m=1}^{\frac{M-1}{2}} (-1)^{m+1} [2\omega]^{-2m-1} Q_{m-1}(\pi) \} e^{-i\omega\pi} \\ &+ \sum_{j=1}^J C_j(m, \omega) e^{-i\omega(\pi - 2\omega_j)} + \sum_{j=J}^1 D_j(m, \omega) e^{i\omega(\pi - 2\omega_j)} \\ &+ \left\{ \frac{1}{2i\omega} [1 + O(\frac{C_\delta}{\omega^{M-1}})] + \sum_{m=1}^{\frac{M-1}{2}} (-1)^m [2\omega]^{-2m} P_{m-1}(\pi) \right. \\ &\left. - i \sum_{m=1}^{\frac{M-1}{2}} (-1)^{m+1} [2\omega]^{-2m-1} Q_{m-1}(\pi) \right\} e^{i\omega\pi}, \quad \omega \notin \mathbf{Z}, \end{aligned} \quad (3.2)$$

in which the $C_j(m, \omega)$ and $D_j(m, \omega)$ can be obtained by comparing (3.2) with (1.8). Besides (3.2), the entire function $y(\pi; \omega)$ is bounded near \mathbf{Z} . Without loss of generality,

we consider the zeros of $Y(\omega) := \omega y(\pi; \omega/i)$ by applying Theorem 2.2 in a suitable strip containing the real axis. We observe the zeros of $Y(\omega)$ spread themselves vertically along the imaginary axis, that is, the zeros of $y(\pi; \omega)$ spread themselves along the real axis. In particular, given the singularity sequence $\{x_{m,k}\}_{m,k}$ which are all distinct by assumption with J elements, we let $\{\omega_j\}_{j=1}^J$ be the rearrangement of $\{x_{m,k}\}_{m,k}$ such that

$$0 = \omega_0 < \omega_1 < \omega_2 < \dots < \omega_J < \omega_{J+1} = \frac{\pi}{2}.$$

We construct following $2J+2$ successive intervals in $[-\pi, \pi]$:

$$L_1 := [-\pi, -\pi + 2\omega_1], L_2 := [-\pi + 2\omega_1, -\pi + 2\omega_2], \dots, L_{J+1} := [-\pi + 2\omega_J, 0], \\ L_{J+2} := [0, \pi - 2\omega_J], \dots, L_{2J+1} := [\pi - 2\omega_2, \pi - 2\omega_1], L_{2J+2} := [\pi - 2\omega_1, \pi].$$

These intervals are applied as the polygons described previously. However, we note that $L_{J+1} \cup L_{J+2}$ combines to generate a sequence of zeros as described by (2.8) and then (2.9) after observing the exponential exponents in (3.2). There are actually $2J+1$ asymptotically rectangular area on duty. Without loss of generality, we take each $\{L_l\}_{l=1}^{2J+2}$ to generate an asymptotically rectangular area $\{R_l\}_{l=1}^{2J+2}$ as described by (2.7) and (2.8). Finally, we note that one can not identify the quantities $\{\mu_{k,j}\}$ in (2.6), because not being able to locate the coefficients $\{m_j\}$ in (2.4) again in (3.2). For our case, the quantities $\{e_k\}$ in (2.5) are equal to 1.

Let zeros in R_l be denoted as z_{n_l} , $l = 1, 2, \dots, 2J+2$ and $n_l \in \mathbb{N}$. Hence, (2.9) implies that

$$(3.3) \quad |N_Y(R_l(\alpha, s, h)) - s(\omega_l - \omega_{l-1})/\pi| < \bar{n}_l - 1 + \epsilon, \quad l = 1, \dots, 2J+2,$$

in which $\bar{n}_l = 2$ by the construction of intervals $\{L_l\}_{l=1}^{2J+2}$ for all l . Here, we define ω_l , $J+1 \leq l \leq 2J+2$ by the symmetry of $\{L_l\}_{l=1}^{2J+2}$. Finally, it is well-known that the zeros of $y(\pi; \omega)$ are simple and real [9], so (3.3) implies

$$z_{n_l} \sim \frac{n_l \pi}{\omega_l - \omega_{l-1}} + O(1), \quad l = 1, \dots, 2J+2; \quad n_l \in \mathbb{N}.$$

Because the zeros of $y(\pi; \omega)$ are symmetric to the imaginary axis, we can rewrite the equation above to be

$$(3.4) \quad z_{n_l} \sim \frac{n_l \pi}{\omega_l - \omega_{l-1}} + O(1), \quad l = 1, \dots, 2J+2; \quad n_l \in \mathbb{Z}.$$

Once again, we note that $\{z_{n_{J+1}}\} \cup \{z_{n_{J+2}}\}$ is the sequence of zeros generated by the interval $L_{J+1} \cup L_{J+2}$. \square

We may observe from (3.3) that the total number of the zeros of $y(\pi; \omega)$ is equal to

$$N_Y\left(\bigcup_{l=1}^{2J+2} R_l(\alpha, s, h)\right) = \sum_{l=1}^{2J+2} N_Y(R_l(\alpha, s, h)) \sim$$

$$\sim \sum_{l=1}^{2J+2} s(\omega_l - \omega_{l-1})/\pi + o(1) = s + o(1),$$

which implies the zeros of $y(\pi; \omega)$ denoted as $\{z_n\}_{n \in \mathbb{Z}}$ have the following asymptotics: $z_n \sim n + O(1)$, $n \in \mathbb{Z}$, which matches with the Counting Lemma in [9, p. 36]. The bounded error term in (3.4) avoids the possible contradiction to the classic asymptotics [9].

СПИСОК ЛИТЕРАТУРЫ

- [1] R. Carlson, R. Threadgill and C. Shubin, Sturm-Liouville problems with finitely many singularities, *Journal of Mathematical Analysis and Applications*, V. 204, 74–101 (1996).
- [2] R. Carlson, A Borg-Levinson theorem for Bessel operators, *Pacific Journal of Mathematics*, Vol. 177, No. 1, 1–26 (1997).
- [3] L. -H Chen, An uniqueness result with some density theorems with interior transmission eigenvalues, *Applicable Analysis*, Vol. 94, No. 8, 1527–1544 (2015).
- [4] D. G. Dickson, Zeros of exponential sums, *Proc. Amer. Math. Soc.*, 16, 84–89 (1965).
- [5] D. G. Dickson, Expansions in Series of Solutions of Linear Difference-Differential and Infinite Order Differential Equations with Constant Coefficients, *Memoirs of the American Mathematical Society*, Rhode Island, USA, No. 23, 1957.
- [6] R. O. Hryniv and Y. V. Mykytyuk, Eigenvalue asymptotics for Sturm-Liouville operators with singular potential, *Journal of Functional Analysis*, 238, 27–57 (2006).
- [7] V. A. Marchenko, *Sturm-Liouville Operators and Applications*, Birkhaeuser Verlag, Basel, 1986.
- [8] B. Ja. Levin, *Distribution of Zeros of Entire Functions*, revised edition, *Translations of Mathematical Monographs*, American Mathematical Society, 1972.
- [9] J. Pöschel and E. Trubowitz, *Inverse Spectral Theory*, Academic Press, Orlando, 1987.

Поступила 20 апреля 2016