

PHYSICAL CONDITIONS IN REGIONS OF LINE FORMATION OF SOLAR TYPE STARS

L.H.RODRÍGUEZ-MERINO¹, O.CARDONA¹, A.FLORES²

Received 8 May 2015

Accepted 24 June 2015

We present the physical parameters of the regions with spectral line formation of a set of stars classified as G2. The method employed in this analysis only requires the full width at half maximum of each spectral line, the effective temperature of the star and few basic atomic parameters. The physical parameters estimated in this work are compared with results obtained with stellar atmosphere models.

Key words: *line:formation (Physical conditions): stars: solar-type*

1. *Introduction.* Solar type stars play an important role in the search of life. Currently there are several instruments in operation with the main goal to look for planets located in habitable zones (e.g. The Kepler Mission). This type of star presents the optimal conditions for sustaining life because of the low ultraviolet emission, therefore it is mandatory to understand as well as possible the physics of their atmospheres.

Several astrophysical groups have developed numerical codes to model both the stellar atmospheres and the line profile spectra, in some cases codes and results are available to the astronomical community, like Atlas9 and Synthe codes built by Kurucz [1,2]¹ or TLUSTY and SYNSPEC codes built by Hubeny & Lanz [3]². The numerical codes compute the atmosphere models based on different arguments; Local Thermodynamics Equilibrium (LTE), Non Local Thermodynamics Equilibrium (NLTE), plane parallel atmospheres, spherical atmospheres, etc. Once a stellar atmosphere model has been computed is possible to calculate the spectral energy distribution emitted by the atmosphere model, within the process of computing the atmosphere model and the spectrum is feasible to follow the change of some physical parameters as function of the physical depth in the atmosphere.

In order to find out an alternative procedure to make a diagnostics of the physical conditions of regions in stellar atmosphere where spectral lines are being formed, Cardona [4] has performed a statistical mechanics analysis of

¹ <http://kurucz.harvard.edu/>

² <http://nova.astro.umd.edu/>

the atomic line broadening by thermal energy fluctuations in a gaseous system. He found that the width of an atomic spectral line is strongly correlated with the physical conditions of the line forming region. The physical parameters associated with the full width at half maximum (fwhm) of a spectroscopic line are the temperature, the number density of particles, and the principal quantum numbers related with the transition which produces the line. Based on this method, Cardona et al. [5] have performed an analysis of the region in the atmosphere of two stars, the white dwarf star G191-B2B and the variable star V803-Cen, where the Lyman- ζ line is being formed. Their conclusions were compared with results of atmosphere models.

Now using the equations obtained by Cardona [4] we determined the physical conditions of regions in the atmosphere of solar type stars where the Balmer and magnesium lines are being formed. In order to be more confident with the parameters estimated in this study we performed a comparison with the results of the solar atmosphere model. The paper is organized as follows: In Section 2 we present the theoretical arguments employed to infer the physical properties of the regions with spectral line formation. In Section 3, we describe the observational data used in this work. The method followed to measure the fwhm of the selected spectral lines, and the results are described in Section 4. In Section 5 we perform a comparison between the newly properties determined with results obtained with the computation of a stellar atmosphere model. The use of the fwhm of an atomic spectral line as a tool to determine physical properties of the stellar atmosphere is discussed in Section 6.

2. Broadening of spectral lines. The spectral lines of an isolated atom are, in principle, nearly perfectly sharp, however due to the finite lifetime of the energy levels, the lines are naturally broaden. The spectral energy distribution emitted by the plasma of a stellar atmosphere is plagued of several absorption lines. The profile of any spectroscopic line is fully related with the physical properties of the forming region. The traditional approach employed to model the profile of the spectral lines is based on considering that there are, basically, three processes which broaden the spectroscopic lines; (1) the natural broadening which is produced by the interaction of the light with the atoms, (2) the pressure broadening which is essentially caused by the collisions between atoms, and (3) the thermal broadening which give rise because of the velocity distribution of the ensemble of atoms. The broadening of the spectroscopic lines produced by any of the processes quoted above is usually described with an absorption coefficient per atom (α). The combined result of the different processes is given by the convolution of all the absorption coefficients,

$$\alpha(\text{total}) = \alpha(\text{natural}) \cdot \alpha(\text{pressure}) \cdot \alpha(\text{thermal}). \quad (1)$$

The absorption coefficients of the natural and pressure broadening are described with Lorentzian profiles, meanwhile the absorption coefficient of the thermal broadening has a Gaussian profile, the convolution of these profiles results in a new profile, the Voigt profile.

An alternative method developed to understand the broadening of the spectroscopic lines was introduced by Cardona [4]. He based his study on the analysis of the energy fluctuations of a thermodynamic system. The energy fluctuations of a system can be produced by several kinds of processes, like the collisions between particles of the ensemble, therefore this alternative analysis must provide us of equations which describe very closely the broadening of spectral lines. Equations given by Cardona [4] (Eqs. (14) and (16) in his work) relate the full width at half maximum of the spectroscopic lines of hydrogenic and nonhydrogenic atoms (respectively) with the physical conditions of the line forming region, and with few basic atomic parameters, obviously some mechanisms of line broadening, like the stellar rotation or the instrumental effects, are not considered by these equations. Although in all stellar spectra most of the spectral lines are in absorption, it is very common to find the line width at half height denoted as the full width at half maximum.

The main difference between the two methods given above is that the first method describes the broadening of spectroscopic lines as function of thermodynamic quantities and quantum mechanical parameters (non-easy to determine), meanwhile the second method describes the broadening of spectral lines mostly based on thermodynamic quantities. Therefore, the equation based on the analysis of energy fluctuations can be used to explore the physical properties of the line forming regions (see Cardona et al. [5]).

3. Observational Data. We have performed the analysis of several absorption lines that are important features in the spectrum of a solar type star. The selected lines are four Balmer lines ($H\alpha$, $H\beta$, $H\gamma$ and $H\delta$), and the lines of the magnesium triplet (Mgb1, Mgb2 and Mgb3). The set of spectra used to study the regions of line formation was retrieved from the ELODIE library V3.1³ through Hyper-Leda database⁴. The archive hosts around two thousand spectra, and it covers the whole spectral type range. The wavelength range covered by any spectrum of the ELODIE Library spans from around 3900 Å to 6800 Å (see Prugniel & Soubiran [6]; Prugniel et al. [7]). It is possible to retrieve fluxes at two resolutions ($R=42000$ and $R=10000$), in case of the spectra at high resolution the fluxes are normalized to the pseudo-continuum. In order to avoid as much as possible the effects of spectral line

³ <http://www.obs.u-bordeaux1.fr/m2a/soubiran/elodielibrary.html>

⁴ <http://leda.univ-lyon1.fr/11/spectrophotometry.html>

blending we have decided to use the spectral energy distributions at high resolution, therefore we have selected the normalized flux of twenty-four solar type stars. We also have used the Solar Flux Atlas (Kurucz et al. [8]) retrieved from the Kurucz web page, which includes the optical part of the solar spectrum measured at high resolution. Table 1 shows the stars classified as G2 in the

Table 1

LIST OF STARS EMPLOYED TO MEASURE THE FWHM OF THE
SELECTED SPECTRAL LINES. FITS FILES OBTAINED FROM
ELODIE LIBRARY PROVIDED THE STELLAR ATMOSPHERE
PARAMETERS

Star	Spectral type ELODIE	Spectral type SIMBAD	T_{eff} K	logg	[Fe/H]	$v \sin i$ km/s
Sun		G2 V	5777	4.44	+0.00	2.0
HD 245	G2 V	G2 V	5772	4.15	-0.59	3.0
HD 3628	G2 V	G2 V Va.	5699	4.02	-0.18	1.0
HD 4307	G2 V	G0 V	5744	3.85	-0.29	2.0
HD 9562	G2 IV	G1 V	5777	3.85	+0.125	4.0
HD 10307	G2 V	G1.5 V	5793	4.11	-0.02	3.0
HD 12235	G2 IV	G2 IV	5858	3.89	+0.180	5.0
HD 12846	G2 V	G2 V	5727	4.41	-0.25	2.0
HD 14802	G2 V	G1 V	5830	3.89	-0.06	4.0
HD 23050	G2 V	G2 V	5772	4.04	-0.43	3.0
HD 76752	G2 V	G2 V	5708	4.23	+0.00	2.0
HD 81809	G2 V	G5 VFe-1CH-0.8	5610	3.83	-0.36	1.8
HD 84737	G2 V	G0.5 Va	5768	3.89	+0.096	2.3
HD 89010	G2 IV	G1.5 IV-V	5669	3.84	-0.01	3.5
HD 105546	G2 III	G2 IIIw	5016	2.21	-1.58	5.2
HD 119550	G2 V	G2 V	5723	3.83	-0.09	4.0
HD 126868	G2 III	G2 IV	5778	3.81	-0.02	14.3
HD 137107	G2 V	G0 V	5919	4.21	-0.11	5.3
HD 143761	G2 V	G0 V	5749	4.02	-0.26	4.0
HD 159181	G2 II	G2 Iab:	5368	1.22	-0.13	11.6
HD 168009	G2 V	G2 V	5728	4.14	-0.04	3.0
HD 186408	G2 V	G1.5 Vb	5732	4.10	+0.108	4.0
HD 209750	G2 Ib	G2 Ib	5399	0.95	-0.06	7.9
HD 221170	G2 IV	G2 IV Va	4561	1.10	-2.10	7.4
HD 225239	G2 V	G2 V	5590	3.75	-0.47	2.0

ELODIE library, it also provides the spectral type, the luminosity class (according to ELODIE library and SIMBAD database), the atmospheric parameters (effective temperature, surface gravity, and metallicity) and the rotational velocity. The atmosphere parameters of the stars from ELODIE library have been retrieved from the FITS file of each spectrum (the set of atmospheric parameters used

were determined based on internal comparisons within the library) and the rotational velocity values are reported by Glebocki & Gnacinski [9].

4. *Analysis of the Line Forming Regions.* All the physical properties imprinted in the spectral energy distribution (SED) emitted by a star are due to physical processes produced in the stellar atmosphere. The continuum arises from regions of the stellar atmosphere where the Rosseland optical depth is $\tau_{\text{ross}} = 2/3$, in this part of the atmosphere the temperature is equal to the effective temperature. Meanwhile absorption lines of the SED are being forming in regions located between the continuum forming region and the top of the atmosphere. Cardona [4] found that the full width at half maximum of an atomic spectral line is given by




Fig.1. The panels display the spectral energy distribution of the Sun around four absorption features; H α , H β , H δ , and Mg b₁. The full width at half maximum of each spectral line is displayed with a discontinuous line, and its value is showed on the right lower part of each panel.

$$w = \frac{2\pi a_0}{3\alpha Z_{eff}} kT \sqrt[3]{N} (n_{ieff}^2 + n_{feff}^2), \quad (2)$$

where N and T are physical parameters of the region of line formation; N is number density of particles, and T is the temperature, n_{ieff} and n_{feff} are the effective principal atomic quantum numbers of the levels producing the line, Z_{eff} is the effective charge of the atom, k and a_0 are the Boltzmann constant and the Bohr radius, respectively, and α is a numerical constant. In order to explore the physical conditions of the regions where the selected spectroscopic lines are being formed we measured the full width at half maximum of the spectral lines.

The selected lines produce prominent absorption features in the spectrum of a solar type star, therefore they have played important roles in several works, e.g. see Worthey & Ottaviani [10] and Worthey et al. [11]. According to




Fig.2. Similar to Fig.1, in this case the star is HD 12846.

ELODIE library all the chosen stars have assigned a G2 spectral type classification, most of them are identified as dwarfs, only few stars are classified as giants or super-giants. However, according to the SIMBAD database, there are some stars that are not classified with a G2 spectral type (see Table 1), then one have to keep in mind that there could be some stars misclassified. Stars with different luminosity class from our selected set of spectra will allow us to explore regions under different physical conditions.

The shape of the absorption line is known as the Voigt profile which is the result of the convolution of Gaussian and Lorentzian profiles (see Section 2). In order to measure the width of a selected spectral feature at the half maximum we performed a linear interpolation of the spectral energy distribution around the minimum of the selected line. The interpolation used a wavelength step of around 0.001 \AA . Because of we have employed the normalized spectra of the ELODIE library, then we only had to measure the width of the interpolated line at the middle distance between the minimum of the flux and the unit. Fig.1 and 2 display parts of the spectrum of the Sun and the star HD 12846, showing the $H\alpha$, $H\beta$, $H\delta$ and the magnesium lines.




Fig.3. In few cases, the spectral line blending does not allow us to measure the fwhm of the selected spectral line. The spectral region close to the $H\delta$ line of the HD 9562 star is populated by several spectral lines making difficult the measurement of the fwhm of this line.

The discontinuous line in each panel describes the full width of the spectroscopic line at the half of the maximum. The fwhm measured is given at the right lower part of each panel. Although we have used a set of spectra with high spectral resolution we could not avoid completely the effects of the line

blending in all the measurements of the fwhm. Fig.3 shows that the spectral region close to the $H\delta$ line of the star HD 9562 is populated by several spectral lines which makes difficult to measure the fwhm of this Balmer line.

The files with normalized fluxes of the ELODIE library do not provide the errors of the stellar fluxes. In order to estimate the error in the measurement of the width of a spectroscopic line we have followed a Montecarlo method. We have selected an spectrum, and we have generated randomly several similar spectra. We used the RMS of the selected spectrum to generate the set of spectra. Using the generated spectra we obtained a set of widths of the

Table 2

FWHM (in \AA) OF THE SELECTED SPECTRAL LINES OF THE SUN AND G2-TYPE STARS. IN FOUR CASES, IT WAS NOT POSSIBLE TO MEASURE THE WIDTH OF THE LINE (see the text)

Star	$H\alpha$	$H\beta$	$H\gamma$	$H\delta$	Mgb3	Mgb2	Mgb1
Sun	1.38	1.19	1.21	1.40	0.71	0.96	1.32
HD 245	1.44	1.09	1.03	1.11	0.56	0.75	0.94
HD 3628	1.36	1.10	1.02	1.08	0.60	0.86	1.25
HD 4307	1.50	1.13	0.96	1.06	0.52	0.66	0.80
HD 9562	1.41	1.20	1.11	-	0.60	0.78	1.18
HD 10307	1.49	1.19	1.11	1.35	0.63	0.83	1.23
HD 12235	1.54	1.27	1.20	1.34	0.60	0.77	1.18
HD 12846	1.37	1.08	0.99	1.06	0.69	1.00	1.35
HD 14802	1.47	1.20	1.20	1.11	0.55	0.65	0.86
HD 23050	1.40	1.10	1.01	1.11	0.54	0.73	0.94
HD 76752	1.33	1.08	1.04	-	0.72	0.95	1.34
HD 81809	1.43	1.08	0.97	1.20	0.64	0.88	1.22
HD 84737	1.52	1.18	1.13	1.20	0.59	0.77	1.11
HD 89010	1.49	1.15	1.01	1.21	0.61	0.78	1.19
HD 105546	1.44	0.94	0.78	0.78	0.40	0.34	0.40
HD 119550	1.48	1.14	0.99	1.05	0.55	0.67	0.97
HD 126868	1.85	1.25	1.03	1.07	0.67	0.79	1.12
HD 137107	1.52	1.29	1.14	1.24	0.58	0.73	1.07
HD 143761	1.42	1.08	1.04	1.29	0.59	0.77	1.02
HD 159181	2.84	1.88	-	1.22	0.71	0.77	1.18
HD 168009	1.36	1.15	1.05	1.25	0.66	0.90	1.28
HD 186408	1.38	1.19	1.08	1.24	0.69	0.94	1.31
HD 209750	2.49	1.64	-	1.11	0.69	0.71	1.14
HD 221170	1.40	0.93	0.72	0.64	0.39	0.32	0.37
HD 225239	1.43	1.02	0.87	1.10	0.52	0.63	0.79

same spectroscopic line. A simple statistical analysis allowed us to determine that the error in the measurement of the fwhm of any spectral line is negligible. Table 2 gives the value of the fwhm of all selected spectral lines. In three cases, HD 9562 ($H\delta$), HD 159181 ($H\gamma$), HD 209750 ($H\gamma$) were not possible to

measure the width of the line due to problems with the line blending, and in the case of the star HD 76752 ($H\delta$) the spectrum shows an emission line in the region of the selected spectral feature.

Fig.4 displays the values of fwhm of the selected spectral lines. The set of values of the fwhm of each spectral line spans around a common value.




Fig.4. The fwhms of each selected spectral line are distributed around a common value. Two special cases are; the Balmer lines, which are formed in the outer parts of the atmosphere, of giant stars are wider, while the magnesium lines of the metal poor stars are thinner (see the text).

The giant and super-giant stars (diamond, triangle and square, respectively) departure from the trend followed by most of the stars because they present stronger Balmer lines. On the other side, stars HD 105546 and HD 221170 (the most metal poor stars of the set) display thinner spectral lines than the common values, in particular their magnesium lines. These departures from the mean values are produced by particular physical phenomena, whose study is outside the scope of this work.

We used Eq. (2) and the full width at half maximum of each selected spectral line to estimate the density of particles of the region where the line is being formed. Fig.5 displays, according to Eq. (2), the change of the fwhm of the $H\alpha$ line as function of the number density of particles and temperature (tilted lines), we employed five temperatures; 100, 90, 80, 70 and 60% of the effective temperature of the Sun, and the horizontal line correspond to the fwhm of the $H\alpha$ line measured in the solar spectrum. This plot allows us to infer the density of particles of the region in the solar atmosphere where the $H\alpha$ line is being formed. Cardona et al. [5] have shown that the Balmer lines as well as the magnesium lines are formed in a region of the stellar atmosphere

where the temperature is close to the 80% of the effective temperature, therefore we have employed Eq. (2), three temperatures; 85, 80 and 75% of the effective temperature of each star, and the full width at half maximum of the selected spectral lines to determine the number density of particles in the regions where the spectroscopic lines are being formed. Table 3 gives the number density of particles of the regions where each selected spectral line is being formed. The density of particles of the region where the $H\alpha$ line is being formed has a common value of $N \sim 12 \times 10^{15}$ particles cm^{-3} . There are four stars; HD 126868, HD 159181, HD 209750 and HD 221170, which the number density of particles estimated is larger than the common value, the reason could be that they are fast rotators, therefore their spectroscopic lines are strongly affected by the rotation. The $H\beta$ absorption lines are less affected by the rotation, and the rest of the analysed Balmer lines do not depart from the respective common value. Also it is important to note that the magnesium lines are good tracers of metallicity, the stars HD 105546 and HD 221170 have the lowest chemical abundances (see Table 1), and probably this is the reason that the number density of their line forming regions have the lowest values.




Fig.5. The full width at half maximum of the $H\alpha$ line as function of the particle density. The tilted lines correspond to percentages of the effective temperature (100, 90, 80, 70 and 60%) of the Sun. The horizontal line shows the width of the $H\alpha$ line measured using the Solar Flux Atlas (Kurucz et al. [8]). The number density of particles where the $H\alpha$ line of the Sun is being formed is around 1×10^{16} particles cm^{-3} .

Table 3

NUMBER DENSITY OF PARTICLES ($\times 10^{15}$ particles cm^{-3}) OF THE REGIONS WHERE THE ABSORPTION LINES ARE BEING FORMED

Star	$N(\text{H}\alpha)$	$N(\text{H}\beta)$	$N(\text{H}\gamma)$	$N(\text{H}\delta)$	$N(\text{Mgb3})$	$N(\text{Mgb2})$	$N(\text{Mgb1})$
Sun	9.96 ± 3.49	10.89 ± 4.17	7.23 ± 1.84	6.12 ± 1.32	24.77 ± 21.59	25.84 ± 23.45	64.10 ± 144.50
HD 245	11.30 ± 4.48	8.15 ± 2.33	4.45 ± 0.70	2.98 ± 0.31	11.64 ± 4.76	11.93 ± 4.99	23.13 ± 18.80
HD 3628	9.89 ± 3.44	8.70 ± 2.67	4.49 ± 0.71	2.33 ± 0.19	14.88 ± 7.80	18.69 ± 12.28	56.50 ± 112.20
HD 4307	12.96 ± 5.91	9.21 ± 2.99	3.66 ± 0.47	2.63 ± 0.24	9.46 ± 3.14	8.25 ± 2.39	14.46 ± 7.35
HD 9562	10.59 ± 3.95	10.85 ± 4.14	5.56 ± 1.09	4.89 ± 0.84	14.28 ± 7.19	13.39 ± 6.28	45.63 ± 73.14
HD 10307	12.39 ± 5.39	10.49 ± 3.86	5.51 ± 1.07	5.31 ± 0.99	16.40 ± 9.44	16.00 ± 8.99	51.25 ± 92.35
HD 12235	13.22 ± 6.13	12.33 ± 5.34	6.73 ± 1.59	5.02 ± 0.88	13.69 ± 6.59	12.34 ± 5.36	43.74 ± 67.26
HD 12846	9.97 ± 3.50	8.12 ± 2.31	4.05 ± 0.58	2.66 ± 0.25	22.30 ± 17.48	28.95 ± 29.47	70.13 ± 172.89
HD 14802	11.67 ± 4.78	10.55 ± 3.92	6.83 ± 1.64	2.89 ± 0.29	10.71 ± 4.03	7.54 ± 2.00	17.19 ± 10.36
HD 23050	10.39 ± 3.79	8.38 ± 2.47	4.20 ± 0.62	2.98 ± 0.31	10.44 ± 3.83	11.00 ± 4.24	23.13 ± 18.80
HD 76752	9.21 ± 2.98	8.20 ± 2.36	4.74 ± 0.79	3.08 ± 0.33	25.59 ± 23.01	25.07 ± 22.06	69.27 ± 168.70
HD 81809	12.05 ± 5.10	8.64 ± 2.63	4.05 ± 0.58	4.10 ± 0.59	18.93 ± 12.60	20.99 ± 15.49	55.06 ± 106.64
HD 84737	13.32 ± 6.26	10.36 ± 3.78	5.89 ± 1.22	3.77 ± 0.50	13.64 ± 6.54	12.94 ± 5.89	38.16 ± 51.18
HD 89010	13.22 ± 6.13	10.10 ± 3.59	4.43 ± 0.69	4.08 ± 0.58	15.89 ± 8.87	14.17 ± 7.03	49.52 ± 86.20
HD 105546	17.23 ± 10.42	7.96 ± 2.23	2.95 ± 0.31	1.58 ± 0.09	6.47 ± 1.47	1.69 ± 0.10	2.71 ± 0.26
HD 119550	12.59 ± 5.58	9.57 ± 3.22	4.06 ± 0.58	2.59 ± 0.24	11.32 ± 4.50	8.73 ± 2.68	26.07 ± 23.89
HD 126868	23.89 ± 20.09	12.25 ± 5.27	4.44 ± 0.69	2.66 ± 0.25	19.88 ± 13.91	13.90 ± 6.80	38.99 ± 53.49
HD 137107	12.33 ± 5.34	12.53 ± 5.53	5.60 ± 1.10	3.85 ± 0.52	11.99 ± 5.05	10.20 ± 3.66	31.63 ± 35.17
HD 143761	10.97 ± 4.24	8.02 ± 2.26	4.64 ± 0.76	4.74 ± 0.79	13.78 ± 6.66	13.06 ± 6.01	29.90 ± 31.47
HD 159181	107.79 ± 408.77	51.99 ± 94.97	5.70 ± 1.14	4.92 ± 0.85	29.50 ± 30.63	16.05 ± 9.08	56.87 ± 113.69
HD 168009	9.74 ± 3.33	9.79 ± 3.37	4.83 ± 0.82	4.36 ± 0.67	19.50 ± 13.36	21.09 ± 15.64	59.75 ± 125.41
HD 186408	10.16 ± 3.63	10.83 ± 4.13	5.24 ± 0.97	4.24 ± 0.63	22.24 ± 17.40	23.99 ± 20.22	63.91 ± 143.66
HD 209750	71.41 ± 179.28	37.11 ± 48.41	6.27 ± 1.38	3.64 ± 0.47	26.61 ± 24.93	12.37 ± 5.39	50.40 ± 89.29
HD 221170	21.05 ± 15.57	10.26 ± 3.71	3.08 ± 0.33	1.16 ± 0.05	7.97 ± 2.23	1.88 ± 0.12	2.86 ± 0.29
HD 225239	12.19 ± 5.22	7.35 ± 1.90	2.95 ± 0.31	3.19 ± 0.36	10.26 ± 3.70	7.78 ± 2.13	15.11 ± 8.05

5. *Comparisons.* In order to show that the results obtained with Eq. (2) are in accordance with physical properties of the stellar atmospheres of G2-type stars we decided to compare these results with physical parameters of the solar atmosphere model. The atmosphere model allows us to know several parameters, like the particle density (N), the temperature (T), the pressure (P), etc., as function of the geometrical depth (Rosseland optical depth, τ_{ross}) inside the atmosphere. It is important to keep in mind that a stellar atmosphere model provides information which describes approximately the formation of spectral lines (see Cardona et al. [5]), however it can be a good benchmark. We decided to calculate the solar atmosphere model using the code developed by Crivellari et al. [12]. The effective temperature used to compute the solar model was 5770 K, the logarithm of the surface gravity was 4.44 dex, and the model has been calculated using the solar abundance. The spectral lines are being formed in regions with $\tau_{\nu} = 1$. Fig.6 shows the Rosseland optical depth as function of the frequency, the model allows us to determine that in the solar atmosphere the $\text{H}\alpha$ absorption

line is being formed around a region where the Rosseland optical depth is $\sim 2.42 \times 10^{-2}$, at this optical depth the temperature is ~ 4600 K and the pressure is ~ 4800 dy cm $^{-2}$, therefore the density of particles is around 7×10^{15} particles cm $^{-3}$. The number density of particles obtained with Eq. (2) has been $9.96 \pm 3.49 \times 10^{15}$ particles cm $^{-3}$, which is very close to the previous result. On




Fig.6. The Rosseland optical depth (τ_{ross}) as function of the frequency given by the solar atmosphere model.

the other side, the atmosphere model lets us know that the H δ absorption line is being formed in a region where the optical depth is around 2.23×10^{-1} , at this depth the temperature is ~ 5300 K and the pressure is ~ 68000 dy cm $^{-2}$, therefore the density of particles is $\sim 90 \times 10^{15}$ particles cm $^{-3}$. Using Eq. (2) we have obtained that the number density of particles is $6.12 \pm 1.32 \times 10^{15}$ particles cm $^{-3}$, which is lower than the previous result. It is important to keep in mind two points; first, the atmosphere model describes approximately the formation of spectral lines, then we do not expect to find a very close matching between both results, and second, the H α absorption line is formed in outer regions of the atmosphere where the temperature is lower, thus large amounts of hydrogen are capable to perform a transition from state $n=2$ to the state $n=3$. While, the H δ line is formed in regions at higher temperatures, thus there should not be large amounts of hydrogen capable to perform a transition from state $n=2$ to state $n=6$.

6. *Discussion and Conclusions.* We have used 25 spectra of stars classified as G2 spectral type in the ELODIE library, and the spectrum of the Sun to measure the full width at half maximum of a set of absorption lines. Most of the stars are dwarfs. The set of measurements of each selected spectral feature ranges around a common value. In the case of the Balmer lines, we found four stars whose fwhm departs from the common values because of they are rapid rotators. We also analysed two stars which show magnesium lines thinner than the common values, the most likely reason is that these stars are metal poor. The small spread of the fwhms around the common values could be due to the spectral misclassification of some stars, blending effects, strong magnetic fields, instrumental broadening, or other phenomena.

We have determined the number density of particles of regions with spectral line formation in the atmosphere of the Sun and in the atmospheres of solar type stars, we used the fwhm of the spectral lines, the effective temperatures of the stars, and few basic atomic parameters to infer the number density of particles. We find out that the number density of particles in regions where the Balmer lines are being formed decreases with the increases of quantum number of the transition, the number density of particles in regions where the $H\alpha$ absorption line is being formed is three times higher than the number density of particles in regions where the $H\delta$ line is being formed. Possibly these results were already expected, however in this work we have used a new method that allows us to quantify directly the changes in the physical conditions of the stellar atmospheres. We also find out that the number density of particles in regions where the magnesium triplet absorption lines are being formed seems to change as function of the stellar metallicity. The magnesium spectral lines have been employed as tracers of the chemical abundances of the stars, then our results seems to follow a similar behaviour. It is important to point out that the equation employed in this study does not take into account broadening of the spectral lines produced by stellar rotation, stellar winds, or other phenomena. Thus, results obtained for the stars HD 126868, HD 159181, HD 209750 and HD 221170 should be taken with cautions.

We have shown that the number density of particles of the regions with spectral line formation estimated in this work is in agreement with the physical properties of the solar atmosphere model. According with the atmosphere model, the $H\alpha$ line is being formed in the outer parts of the atmosphere where many more hydrogen atoms are not ionized, therefore most of these atoms can carry out the transition from state $n=2$ to the state $n=3$, then the densities of particles of the region where the $H\alpha$ absorption line is being formed obtained with both methods are very similar. On the other hand, the fwhm of the $H\delta$ spectroscopic line allowed us to infer a number density of particles of the regions where the line is being formed which is lower than the density

estimated with the atmosphere model. In spite that this result seems to be wrong, it is fine, because the $H\delta$ line is being formed at deeper regions, so the temperature is higher, therefore the number of particles performing the transition from state $n=2$ to the state $n=6$ is lower. Thus the results presented in this work are in agreement with the expected physical properties in the atmospheres of G2 spectral type stars.

The full width at half maximum of an absorption line is a useful tool to study the physical conditions of the region where the spectral line is being formed. It allows us to infer physical properties of the plasma in the atmosphere of a star that the atmosphere model can not do.

This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France.

¹ Instituto Nacional de Astrofísica, Óptica y Electrónica, Tonantzintla, México, e-mail: lino@inaoep.mx

² Facultad de Ingeniería, Universidad Autónoma del Carmen, Ciudad del Carmen, México, e-mail: aflores@pampano.unacar.mx

ФИЗИЧЕСКИЕ УСЛОВИЯ В ОБЛАСТЯХ ОБРАЗОВАНИЯ ЛИНИЙ ЗВЕЗД СОЛНЕЧНОГО ТИПА

Л.РОДРИГЕЗ-MERINO¹, О.КАРДОНА¹, А.ФЛОРЕС²

Представляем физические параметры областей со спектральными линиями образования ряда звезд, классифицированных как G2. Используемый в этом анализе метод требует только полную ширину при половине максимума каждой спектральной линии, эффективную температуру звезды и несколько атомных параметров. Физические параметры, оцененные в этой работе, сравнены с результатами, полученными с моделями звездных атмосфер.

Ключевые слова: *линии: образование (физические условия): звезды солнечного типа*

REFERENCES

1. *R.Kurucz*, ATLAS9 Stellar Atmosphere Programs and 2 km/s grid. Kurucz CD-ROM No. 13. Cambridge, Mass.: Smithsonian Astrophysical Observatory, 1993a, 13.
2. *R.Kurucz*, SYNTHE Spectrum Synthesis Programs and Line Data. Kurucz CD-ROM No. 18. Cambridge, Mass.: Smithsonian Astrophysical Observatory, 1993b, 18.
3. *I.Hubeny*, *T.Lanz*, *Astrophys. J.*, **439**, 875, 1995.
4. *O.Cardona*, *Astrophysics*, **54**, 75, 2011a.
5. *O.Cardona*, *A.Flores*, *L.H.Rodrigues-Merino*, *Astrophysics*, **54**, 538, 2011b.
6. *P.Prugniel*, *C.Soubiran*, *Astron. Astrophys.*, **369**, 1048, 2001.
7. *P.Prugniel*, *C.Soubiran*, *M.Koleva*, *D. Le Borgne*, arXiv:astro-ph/0703658, 2007.
8. *R.L.Kurucz*, *I.Furenlid*, *J.Brault*, *L.Testerman*, National Solar Observatory Atlas, Sunspot, New Mexico: National Solar Observatory, 1984.
9. *R.Glebocki*, *P.Gnacinski*, *VizieR Online Data Catalog*, 3244, 2005.
10. *G.Worthey*, *D.L.Ottaviani*, *Astrophys. J. Suppl. Ser.*, **111**, 377, 1997.
11. *G.Worthey*, *S.M.Faber*, *J.J.Gonzalez*, *D.Burstein*, *Astrophys. J. Suppl. Ser.*, **94**, 687, 1994.
12. *L.Crivellari*, *O.Cardona*, *E.Simonneau*, *Astrophysics*, **45**, 480, 2002.

