



UDC 663.222:663.252.41

Investigating the Impact of Yeasts Additives on the Quality of the Red Wines: Autolyzes of Yeast and Yeast Derivatives Introduced During Alcoholic Fermentation

K.N. Kazumyan, M.N. Mikayelyan, E.R. Gevorgyan, A.A. Jraghatspanyan

Armenian National Agrarian University

kkazumyan@yahoo.com, mikayelyan.m@mail.ru, elyanora.gevorgyan@mail.ru, artem.jraghatspanyan@mail.ru

ARTICLE INFO

Keywords:

active dry yeast,
maceration,
phenolic compounds,
red grape varieties,
wine color,
wine tasting,
yeast derivatives

ABSTRACT

The second part of the article series is about the study of the influence of yeast and autolyzed yeast derivative on red wines' qualitative and quantitative characteristics. Wine samples in which the preparation process was explained in the first part of the article were investigated. Wine coloring and phenolic compounds were measured. Organoleptic properties of the wines are presented as spider graphs with tasting notes.

Introduction

One of the main quality parameters of red wine is color and phenolic compounds. As mentioned in the study's first article (Kazumyan and Mikayelyan, 2022), Armenian grape varieties have been chosen: Tigrani, Karmrahyut, and Charentsi which are selective grapes with red-colored pulp (juice). These grape varieties have been slightly investigated and our research will be useful for many Armenian winemaking companies. Winemakers use these varieties to enhance the color and taste of red wines. Many winemaking companies refuse to use the Karmratut grape because it has unstable color and coloring matter sediment. Phenolic compounds in red wines also influence the wines' quality, color stability, and aging potential.

Materials and methods

The OIV and EAEU GOST methods were used to assess wine physicochemical indicators. Phenolic compounds are measured using the photocolormetry method (Fracassetti and Gabrielli, 2017; Anikina, 2019; Gerzhikova, 2009; Jacobson, 2006; Ljiljana, 2016; Marnie, 2013; Moreno, 2012).

Armenian wine industry experts at the EVN Yerevan Wine Academy at Armenian National Agrarian University evaluated the organoleptic characteristics of the wines at the tasting room of the EVN Yerevan Wine Academy in collaboration with www.winenet.am/en/, the online resource.

Results and discussion

The corresponding spider diagrams (Aroma wheels) were drawn up based on the tasting results. The sensory analysis of wine quality involves assessing how the wine tastes, smells and looks (Marnie, 2013). The fermentation and aging processes first determine a wine's physicochemical




Figure 1. An aroma wheel and tasting graph for wines made from Tigrani, Karmrahyut, and Charentsi grape varieties (composed by the authors).

composition. According to the matrix principle, sensory evaluation was conducted to determine the relationship between organoleptic assessment and physicochemical indicators of young wine composition. A total of 11 indicators were used to assess aroma perception, five indicators to assess taste perception, and four indicators to assess general perception. Based on all the information obtained, the spider graph of the wine-tasting matrix reflects the aroma wheels (Figure 1 and Table 1). It can be seen that Karmrahyut wine has the strongest flower smell at 3.2, Tigrani wine has the strongest smell of red and black fruits inherent in red wines in general, Karmrahyut wine has a spicy smell at 2.3, yeast smell is about the same as 1.7-1.8; the oak smell is lower in Karmrahyut wine, at 1.4.

The characteristic of taste according to the results of tasting is distributed sweet, most pronounced in Tigrani wine – 2.2. The yeast race used has the property of increased accumulation of glycerol (up to 8 g/l); acidic taste prevails in Karmrahyut and Charentsi wines – 2.5-2.4; tannin taste is expressed in all samples and prevails in Charentsi wine 3.0 (Vidal et al., 2004); the taste of alcohol is expressed in Tigrani wine – 3.4; the taste of bitterness manifests itself in Charentsi wine – 2.3.

The general impression of the harmony taste, according to the characteristics of Tigrani (2.2) is noted below; against 2.6 in Karmrahyut and Charentsi wines, in terms of complexity the Tigrani is inferior to 2.4; in Karmrahyut and Charentsi, respectively 2.7 and 2.6.

Table 1. The wine-tasting results are drawn from Tigrani, Karmrahyut and Charentsi grapes*

Odour							
	Karmrahyut	Charentsi	Tigrani		Karmrahyut	Charentsi	Tigrani
Floral	3.2	2.7	1.8	Dried Fruit	2.4	2.0	2.7
Citrus	2.0	1.6	1.4	Herbaceous	2.0	1.9	2.1
Stone fruit	1.7	1.8	1.9	Spicy	2.3	1.7	1.7
Tropical fruit	1.5	1.6	1.6	Yeast	1.8	1.7	1.7
Red fruit	2.6	2.6	2.4	Oak	1.4	1.7	1.7
Black fruit	2.8	2.7	3.1				
Taste				Overall Impression			
Sweetness	1.8	1.7	2.2	Harmony	2.6	2.6	2.2
Acidity	2.5	2.4	1.8	Complexity	2.7	2.6	2.4
Tannin	2.9	3.0	2.5	Body	2.7	2.9	2.6
Alcohol	2.4	2.8	3.4	Potential	2.6	2.4	2.6
Bitterness	1.7	2.3	1.7	max = 5			

*Composed by the authors.

According to the characteristics of the body of the wine, the prevailing wine sample is Charentsi 2.9. The development potential is estimated at 2.6 for Tigrani and Karmrahyut wines and 2.4 for Charentsi wine.

Phenolic compounds play a significant role in plant metabolism. Grapes and wine contain phenolic compounds and their balanced complex determines red wine quality. Grape compounds, especially flavonoids, are significant in high-quality red wines. Flavonoids are mainly concentrated in the solid parts of the grape (Ribereau-Gayon, et al., 2006; Ribereau-Gayon, et al., 2006).

Processing of grapes and the subsequent alcoholic fermentation: after maceration of the grape pomace, the must is enriched with these valuable substances from the solid parts of the grape. According to the degree of oxidation, flavonoids are divided into groups. The mutual transformation of one group into another is possible through oxidation and reduction processes.

Tannins or proanthocyanidins (polymers of flavan-3-ol) are the most common phenols in grapes. Red wines are astringent due to tannins extracted from the skin and the soft parenchyma of the seed (Vidal, 2004; Yakimchenko, 2012). The diffusion coefficient, i.e. the rate at which tannin is extracted into fermenting must, depends on the size of the tannin molecule. Grapes contain monomers, dimers, trimers,

oligomers, and polymers. The degree of polymerization reaches significant levels, and the molecular weight is up to 3500. These are catechin (condensed) tannins. Seed tannins consist of catechin and epicatechin. The degree of their polymerization is 10 units. Tannins from skins and seeds are extracted at different speeds. Extraction from the seeds is more intensive, easier, and in greater quantities than from the skins (Ribereau-Gayon, 2006). During alcoholic fermentation, tannin accumulation peaks on the 7-10th day (Kazumov, 2013). Tannins are easily dissolved in alcohol. Therefore, during alcoholic fermentation with the accumulation of ethyl alcohol, the dissolving ability must increase tannins (Yanniotis, 2007). After fermentation of sugars and enzymatic transformations, tannins polymerize, yielding macromolecules of proteins and polysaccharides. This forms complex compounds that become colloidal and precipitate. The tannins of grape skin and seeds have different effects on the wine taste (Busse-Valverde, 2011). Seed tannins define the wine's structure and body, while skin tannins make the wine soft and smooth (Marnie, 2013).

Catechins, tannins, and anthocyanins are the most concentrated natural antioxidants (Maleti, 2009). Tannins are the first oxidized substance of wine. Other compounds also can be involved in oxidative reactions (Mazza, 1995; Vidal, 2004; Yakimchenko, 2012).

Table 2. Tigrani, Karmrahyut, and Charentsi wine phenolic compounds and color composition*

№	Phenolic compounds				
	Parameters	unit	Charentsi	Karmrahyut	Tigrani
3.1.	Total Flavonoids	mg/l	2263.96	2261.88	1569.72
3.2.	Total anthocyanins	mg/l	368.73	508.45	658.30
3.3.	Total phenols (Gallic acid)	mg/l	2808.93	2740.71	2019.54
3.4.	Folin-Cicalteu Index		64	62	46
4.	Chromatic characteristics				
4.1.	A-420		0.37	0.35	0.45
4.2.	A-520		0.42	0.45	0.57
4.3.	A-620		0.11	0.12	0.19
4.4.	Color Intensity		8.96	9.17	12.19
4.5.	Color Shade		0.88	0.77	0.79
4.6.	Color composition, %				
4.6.1.	A-420 yellow	%	41.22	37.76	37.17
4.6.2.	A-520 red	%	46.79	49.00	46.99
4.6.3.	A-620 blue	%	13.63	15.25	18.82

*Composed by the authors.

Using the results of our tests shown in (Table 2), we determined the total amount of anthocyanins, flavonoids, and phenolic compounds in wine. These compounds are significant for evaluating tannin-anthocyanin reserves to prepare high-quality red wines.

The total amount of phenolic compounds is in the range of 2019.54 mg/l in Tigrani (the minimum amount), in Karmrahyut: 2740.71 mg/l, in Charentsi: 2808.93 mg/l (maximum amount). Total flavonoids – condensed phenolic substances recorded in samples: Tigrani – 1569.72 mg/l or 77.73 % (of the content); Karmrahyut – 2261.88 mg/l or 82.53 %; Charentsi – 2263.96 mg/l or 80.6%. Folin & Ciocalteu's is an international index of total phenolic compound content in wines (International Organisation of Vine and Wine, 2021). The index is for Charentsi – 64, the highest and Tigrani – 46, the smallest.

A grape's anthocyanins determine the color of its future wine (Monagas, M. 2009; Wrolstad, 2005; Burns, 2002). Winemaking using the red method extracts the maximum amount of color pigments from the grape skin during the maceration process in the ongoing alcoholic fermentation and their subsequent preservation in wine (Ribereau-Gayon, 2006). Anthocyanins are polyphenols present in grapes and wine in the form of glycosides and diglycosides. Depending on the grape variety, the anthocyanin composition and total amount in wine differ. The total amount of anthocyanins was recorded as 368.73 mg/l in Charentsi, the lowest amount, 508.45 mg/l in Karmrahyut, and 658.30 mg/l in Tigrani, the highest.

Grapes contain anthocyanins, which cause wine's red color. Additionally, they are found in the thick epidermal layer of the skin. They absorb light and represent many colors ranging from red to blue (Anikina, 2012; Ribereau-Gayon, 2006). The color of young wine obtained through maceration depends on the minerals present in the wort. Anthocyanins (anthocyanidins) are associated with sugar molecules. Externally, anthocyanins are free in grapes and wine and can be linked by ester bonds with acetic and coumaric acids, with substituted hydroxyl and individual groups. The color and shade of anthocyanins are determined by their qualitative and quantitative states. The color composition in the samples was determined on a UNICO 2802 UV/VIS photoelectric colorimeter at 420, 520, and 620 nm, in a 1 mm thick cuvette (International Organisation of Vine and Wine, 2021; Heredia, 1998). As a result of the research, color intensity was obtained: the largest in Tigrani – 12.19; Karmrahyut – 9.17; the smallest in Charentsi – 8.96.

The highest color shade indicator was recorded in the Charentsi sample – 0.88, and in the Karmrahyut and Tigrani samples, respectively – 0.77 and 0.79.

There are three colors in the color composition, each determined by wavelengths of 420 nm (yellow), 520 nm (red), and 620 nm (blue). With over 65 % of the samples being red and blue, the Tigrani sample has the highest combination of these colors. As a result, the color intensity index is 12.19.

Conclusion

Based on the results of this study, we can conclude that the selected active dry yeast and yeast derivatives are beneficial for the production of red wine. Wines made from investigated grapes meet all quality requirements. Based on the tasting results, all wines display a body, typically red and black fruit aromas.

While the wine samples are in their middle stage of complexity and harmony, they were bottled in the spring of 2021 without being seasoned. The wine samples have excellent anthocyanins and the phenolic content is high. The highest total anthocyanins content was found in the Tigrani wine sample and the greatest total phenols were found in the Charenci wine sample.

About 50 % of the total wine color was determined in the Karmrahyut wine sample. The aging potential of red wine is crucial for evaluating it. For reliable and valid research findings, this aspect must be assessed accurately, as well as yeast derivatives should be considered. Following a substantial period of bottle maturation, we plan to perform an aging potential analysis. Typically, this process takes several years.

References

1. Anikina, N.S., Cherviakov, S.N., Gnilomedova, N.V. (2019). Methods for evaluating the color of wines, "The review Analytics and Control", vol. 23, N. 2, - pp. 158-167.
2. Burns, J., Mullen, W., Landrault, N., Teissedre, P.L., Lean, M.E.J., Crozier, A. (2002). "Variations in the profile and content of anthocyanins in wines made from Cabernet Sauvignon and hybrid grapes". J. Agric. Food Chem. 50, - pp. 4096–4102 <https://doi.org/10.1021/jf011233s>.
3. Busse-Valverde, N., Gómez-Plaza, E., López-Roc, J.M., Gil-Muñoz, R., Bautista-Ortín, A.B. (2011). The extraction of anthocyanins and proanthocyanidins from grapes to wine during fermentative maceration is affected by the enological technique. J. Agric. Food Chem. 59, pp.5450–5455 <https://doi.org/10.1021/jf2002188>.

4. Daniela Fracassetti, Mario Gabrielli, Onofrio Corona, Antonio Tirelli, Characterization of Vernaccia Nera (*Vitis vinifera* L.) Grape and Wine, South African Journal of Enology & Viticulture. (2017). Vol. 38, No. 1, - pp. 78-81 <https://doi.org/10.21548/38-1-867>.
5. Gerzhikova, V.G. (2009). Methods of technological control in winemaking, second edition, Simferopol, - p. 304.
6. Heredia, F.J., Francia-Aricha, E.M., Rivas-Gonzalo, J.C., Vicario, I.M., Santos-Buelga, C. (1998). Chromatic characterization of anthocyanins from red grapes - I. pH effect. Food Chem. 63, - pp. 491-498 [https://doi.org/10.1016/s0308-8146\(98\)00051-x](https://doi.org/10.1016/s0308-8146(98)00051-x).
7. International Organisation of Vine and Wine, "Compendium of International Methods of Wine and Must Analysis", OIV-18 RUE D'AGUESSEAU - 75008 PARIS, Edition 2016, Volume 2, - p. 907.
8. International Organisation of Vine and Wine, "Compendium of International Methods of Wine and Must Analysis", OIV-18 RUE D'AGUESSEAU - 75008 PARIS, Edition (2021) Volume 1, - p. 607.
9. Jacobson, J. L. (2006). Introduction to Wine Laboratory Practices and Procedures", United States of America, Springer Science + Business Media, Inc., - p. 375.
10. Kazumov, N.B., Kazumyan, K.N., Sukoyan, M.R. (2013). The technology of Armenian grape, fruit, and berry wines (3rd revised edition). Ancestor, - p. 340.
11. Kazumyan, K.N., Mikayelyan, M.N., Gevorgyan, E.R., Jraghatspanyan, A.A. (2022). "Investigating the Effect of Yeasts and their Derivatives on the Qualitative Indices of Red Wine", Food Science and Technology ANAU, N2/78, - pp.196-201 <https://doi.org/10.52276/25792822-2022.2-196>.
12. Ljiljana, M.B., Dejan, M.G., Ranko, S. (2016). Spectrophotometric characterization of red wine color from the vineyard region of Metohia. Journal of Agricultural Sciences, Belgrade. 61, - pp. 281-290. DOI: 10.2298/JAS1603281B.
13. Maleti, E., Karoglan, J.K., Preiner, D., Claus-Dieter, J.P., Dietrich, H. (2009). Anthocyanin profile and antioxidative capacity of some autochthonous Croatian red wines; Journal of Food, Agriculture & Environment Vol.7 (1), - pp. 48-51.
14. Marnie, O. (2013). Wine: A Tasting Course: Every Class in a Glass, Dorling Kindersley Publishing, Incorporated, - p. 256.
15. Mazza, G., Francis, F.J. (1995). Anthocyanins in grapes and grape products. Crit. Rev. Food Sci. Nutr. 35, pp. 341-371 <https://doi.org/10.1080/10408399509527704>.
16. Monagas, M., Bartolomé, B. (2009). Anthocyanins and anthocyanin-derived compounds. In Wine Chemistry and Biochemistry, 1st ed.; Victoria Moreno Arribas, M., Carmen Polo, M., Eds.; Springer Science+Business Media, LLC: New York, NY, USA, - pp. 439-462 https://doi.org/10.1007/978-0-387-74118-5_21.
17. Moreno, J., Peinado, R. (2012). Enological Chemistry, USA, Elsevier Inc., - p. 429.
18. Ribereau-Gayon, P., Gloires, Y., Maujean, A. Dubourdieu, D. (2006). Handbook of Enology, Volume 2: The Chemistry of Wine and Stabilization and Treatments. John Wiley & Sons, - p. 442 <https://doi.org/10.1002/0470010398>.
19. Ribereau-Gayon, P., D. Dubourdieu. Handbook of Enology, Volume 1, The Microbiology of Wine and Vinifications, 2nd Edition echeand, John Wiley & Sons, 2006, - p. 498 <https://doi.org/10.1002/0470010363>.
20. Tikhonova, A.N., Stribizheva, L.I., Yezhov, E.V., Kachaeva, N.Y. (2011). Influence of Strains of Active Dry Yeast on Organoleptic of Fault, "Winemaking and viticulture", N: 2, - pp. 14-15.
21. Vidal, S., Francis, L., Noble, A., Kwiatkowski, M., Cheynier, V., Waters, E. (2004). Taste and mouth-feel properties of different types of tannin-like polyphenolic compounds and anthocyanins in wine. Anal. Chim. Acta, 513, - pp. 57-65 <https://doi.org/10.1016/j.aca.2003.10.017>.
22. Wrolstad, R.E., Durst, R.W., Lee, J. (2005). Tracking color and pigment changes in anthocyanin products. Trends Food Sci. Tech. 16, - pp. 423-428 <https://doi.org/10.1016/j.tifs.2005.03.019>.
23. Yakimchenko, E.N., Guguchkina, T.I., Belyakova, E.A., Romanishin, P.E. (2012). Phenolic Complex of Canteens and Special Wine from Native Grapes from Don, "Winemaking and viticulture", N: 5, - pp. 27-29.
24. Yanniotis, S., Kotseridis, G., Orfanidou, A., Petraki A. (2007). Effect of ethanol, dry extract, and glycerol on the viscosity of wine, Journal of Food Engineering N: 81, - pp. 399-403 <https://doi.org/10.1016/j.jfoodeng.2006.11.014>.

Accepted on 17.04.2023

Reviewed on 12.06.2023